




YaraVita™
REVIG™



NOTICE

Before using this product, read the entire label

REVIG™

3-0-34

Guaranteed analysis:

Total Nitrogen (N).....**3%**

3% Urea Nitrogen

Soluble Potash (K₂O).....**34%**

Derived from Urea, Potassium formate

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

**2½ gallons/
9.4 litres**

**Net weight
30lb 6oz/
13.8 kg**

F1673

Contents made in Great Britain

Guaranteed & Manufactured by Yara UK Ltd.,

Manor Place, Wellington Road, The Industrial Estate,

Pocklington, York YO42 1DN, U.K.

Manufactured for **Yara North America, Inc.** 100 North Tampa Street,
Suite 3200, Tampa, FL 33602

Customer Service: 800-234-9376

YaraVita REVIG™

APPLICATION RATES AND TIMINGS

APPLY ONLY AS DIRECTED. FOR PROFESSIONAL USE ONLY.

Apples: 1 quart/acre at bud burst. Also, two to three applications of 3 pints/acre at 10 to 14 day intervals starting at petal fall. A further application of 1 quart/acre may also be applied post-harvest before leaf fall. Water rate: 50 gallons/acre.

Apricot, Cherry, Nectarines, Peach: 3 applications of 1 to 2 quarts/acre from stone hardening, with repeat applications at 10 to 14 day intervals. Water rate: 50 to 100 gallons/acre.

Broccoli, Brussel sprouts, Cabbage, Calabrese, Cauliflower: 2 quarts/acre at the 4 to 6 leaf stage. Repeat as required for moderate to severe deficiency at 7 to 14 day intervals. Water rate: 20 gallons/acre.

Canola: 2 quarts/acre at the 4 to 6 leaf stage. Repeat as required for moderate to severe deficiency at 7 to 14 day intervals. Water rate: 20 gallons/acre.

Cereals: 2 quarts/acre from mid-tillering to 2nd node detectable (Zadoks G.S. 25 to 32). Repeat at 10 to 14 day intervals between these growth stages as necessary. Water rate: 20 gallons/acre.

Corn: 2 quarts/acre at the 4 to 8 leaf stage. Water rate: 20 gallons/acre.

Cotton: 2 quarts/acre at squaring and at flowering. Water rate: 20 gallons/acre.

Cucumber, Squash, Water melon, Zucchini (Field Growth): Apply 1 to 2 quarts/acre at 10 day intervals from setting of first fruit/first fruit development. Water rate: 50 gallons/acre.

Grapevines: 1 to 2 quarts/acre at fruit set, pea-sized berries and first color softening/one month before harvest. Water rate: 20 to 50 gallons/acre.

Melons (Field Growth): 2 quarts/acre applied from first fruit development. Repeat at 10 day intervals. Water rate: 50 gallons/acre.

DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES. DO NOT APPLY IF HEAVY RAIN IS EXPECTED. APPLY THIS PRODUCT ONLY TO YOUR LAWN/GARDEN, AND SWEEP ANY PRODUCT THAT LANDS ON THE DRIVEWAY, SIDEWALK, OR STREET, BACK ONTO YOUR LAWN/GARDEN

DIRECTIONS FOR USE

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user. Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use.

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures exceeding 100°F) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request. Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Nuts (Deciduous): 2 quarts/acre at bud break in spring. Water rate: 50 gallons/acre.

Nursery stock/ornamentals: 1 - 2 qts. Per 100 gallon of water. Repeat at 10 - 14 day intervals as necessary. Do not apply during flowering.

Onions: 2 quarts/acre when sufficient leaf area to intercept spray. Water rate: 20 to 50 gallons/acre.

Potatoes: Two applications of 2 quarts/acre during tuber bulking (as soon as first formed tubers are half an inch in diameter). Allow 10 to 14 days between applications. Water rate: 20 gallons/acre.

Raspberry, Strawberry: (Field Growth): 2 quarts/acre at green bud. Water rate: 50 gallons/acre.

Soybeans: 1 quart/acre when the crop is 4 to 6 inches tall. Water rate: 3 to 20 gallons/acre.

Sugar Cane: 2 quarts/acre when cane is between 12 and 50 inches tall. Repeat applications may be necessary at 10 to 14 day intervals. Water rate: 3 to 20 gallons/acre.

Sweet Potatoes: 2 quarts/acre one week after 100% emergence or transplanting. Also, apply at the same rate following recommendation from analysis. Water rate: 20 gallons/acre.

Tobacco: Three applications of 1 to 2 quarts/acre two to three weeks after transplanting (3 to 4 leaf stage) with 10 days between applications. Water rate: 3 to 50 gallons/acre

Tomatoes (Field Growth): 1 quart/acre applied at first truss, fruit set and repeated at 10 day intervals. Water rate: 3 to 50 gallons/acre.

Turfgrass: Apply 1 to 3 fl oz per 1000 sq ft in a water volume of 1.5 Gallons/1000 sq ft every 7 - 14 days as needed during the growing season based on nutritional requirements.

Extended recommendations available at <https://www.yara.us/crop-nutrition/fertilizer-products/yaravita/>

www.yara.us/crop-nutrition/fertilizer-products/yaravita/



WARNING: Cancer and Reproductive Harm – www.P65Warnings.ca.gov