



YaraAmplix™

OPTITRAC™



Net Content
2 1/2 gallons/
9.4 litres
Net weight
24lb 6oz/
11.1 kg

NOTICE

Before using this product, read the entire label

OPTITRAC™

5-0-2

A concentrated liquid fertilizer with micronutrients.

Guaranteed analysis:

Total Nitrogen (N)	5%
5% Urea Nitrogen	
Soluble Potash (K₂O)	2%
Boron (B)	1.1%
Zinc (Zn)	1.1%
1.1% Water Soluble Zinc (Zn)	

Derived from urea, potassium citrate, boron ethanolamine, zinc sulfate, kelp extract.

Density: 9.76 lbs/gal @ 68F

This product is recommended for use in conjunction with a complete fertilization program based on soil or tissue analysis.

F1106

Guaranteed by **Yara North America, Inc.**

100 North Tampa Street, Suite 3800, Tampa, FL 33602

Customer Service: 800-234-9376

Contents made in Great Britain.

Manufactured by **Yara UK Ltd.**, Manor Place, The Industrial Estate, Pocklington, York, YO42 1DN, U.K.

YaraAmplix™ OPTITRAC™

WARNING

This product contains boron which may be injurious to certain crops.
Contact your county agent or field consultant for specific information.

ONLY APPLY TO THE INTENDED APPLICATION SITE. DO NOT APPLY NEAR WATER, STORM DRAINS, OR DRAINAGE DITCHES OR TO ANY IMPERVIOUS SURFACE. DO NOT APPLY IF HEAVY RAIN IS EXPECTED.

APPLICATION RATES AND TIMINGS

Corn: 1-2 pints/acre at the 4 to 8 leaf stage. Water rate: 20 gallons/acre minimum.

Grapes (table): 1-3 pints/acre applied: pre-flowering at 2-6" cluster stretch; post bloom at/after berry set and also during berry sizing. Water rate: 20 gallons/acre minimum.

Grapes (wine): 1-3 pints/acre applied: during 1-4" early shoot growth and again at 10-12" shoot growth when clusters are visible. Avoid applications pre-bloom on varieties that tend to over set; conversely on varieties that over shatter, apply up to 7 days before bloom; and apply post bloom at BB size berries (2-3mm). Water rate: 20 gallons/acre minimum

Lettuce: Field Grown: Three applications of 1-3 pints/acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 gallons/acre minimum.

Melon: 1-3 pints/acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage, pre flower and after fruit set during sizing. Water rate: 20 gallons/acre minimum.

Nuts (deciduous): Apply 1-3 pints/acre at the start of flowering, after nut set during sizing and in season during periods of environmental stress. Also, post harvest applications before leaf senescence. Water rate: 20 gallons/acre minimum.

Potatoes: 1-2 pints/acre applied 7 to 14 days after 100 percent emergence, 2-3 weeks after tuber initiation and (following petiole analysis), during tuber bulking. Water rate: 20 gallons/acre, minimum.

Soya bean: Three applications of 1-3 pints/acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 gallons/acre.


Tomato (Field Grown): Three applications of 0.5 pints/acre at 10 to 14 day intervals commencing at the 4 to 6 leaf stage. Water rate: 20 gallons/acre minimum.

Continuous Cropping Field Grown Varieties: 0.5 pints/acre at the 4 to 6 leaf stage. Applications may be repeated at 10 to 14 day intervals, as necessary. Water rate: 20 gallons/acre.



Extended recommendations available at www.yara.us/crop-nutrition/fertilizer-products/yaravita

YaraAmplix™ OPTITRAC™



9

UN3082

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.
(Zinc sulphate)

Danger

H318: Causes serious eye damage. H315: Causes skin irritation. H411: Toxic to aquatic life with long lasting effects. P280-d: Wear protective gloves/clothing and eye/face protection. P261: Avoid breathing gas, vapor or spray.

P305+P351+P338+P310: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

P302+P352+P333 + P313: IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs, seek medical advice/attention. P391: Collect spillage.
Yara North America, Inc. 100 North Tampa Street, Suite 3800, Tampa, FL 33602
Emergency Tel: Chemtrec 24-hours Emergency Response: 1-800-424-9300

YaraAmplix™ OPTITRAC™

DIRECTIONS FOR USE.

ALWAYS FOLLOW THESE LABEL INSTRUCTIONS

Mix the product thoroughly before use. Slowly add the product to the tank, via the induction hopper, whilst agitating. Top up with water and continue to agitate until spraying is completed.

TANK MIXING/CO-APPLICATION

Read ALL labels carefully and adhere strictly to the instructions for use. Co-application is entirely at the risk of the end-user.

Before co-application of products you, or your advisor, must visit www.tankmix.com for important information. Alternatively, contact Yara for specific advice.

PRECAUTIONS

Only to be used on protected crops if specifically listed. Avoid application under extremes of climate, e.g. rapid drying conditions, extremely slow drying conditions, frost, rain, or when frost or rain are anticipated. Wear suitable gloves and face shield when handling this product. Clean all equipment before and after use. Please refer to the SDS for further information, available at www.yara.com

STORAGE

Keep in a cool, dry chemical store, out of reach of children and livestock. Protect from frost and other climatic extremes. (temperatures $\leq 23^{\circ}\text{F}$ and $> 95^{\circ}\text{F}$) Dispose of contents and container in accordance with all local, regional, national and international regulations.

CONDITIONS OF SALE

This Product is sold subject to the seller's Terms and Conditions of Sale, which are available upon request.

Use of the Product is acceptance by the buyer of the Terms and Conditions of Sale.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>



WARNING: Cancer and Reproductive Harm –www.P65Warnings.ca.gov